

Arrival of extremely volatile-rich high-Mg magmas changes explosivity of Mount Etna

Vadim S. Kamenetsky*

Australian Research Council Centre of Excellence in Ore Deposits and School of Earth Sciences,
University of Tasmania, Hobart, Tasmania 7001, Australia

Massimo Pompilio

Istituto Nazionale di Geofisica e Vulcanologia–Sezione di Pisa, Via della Faggiola 32 Pisa 56126, Italy

Nicole Métrich

Laboratoire Pierre Sue, Centre National de la Recherche Scientifique–Commissariat à l'Energie Atomique,
CE-Saclay, 91191 Gif sur Yvette, France

Alexander V. Sobolev

Max-Planck-Institut für Chemie, Postfach 3060, 55020 Mainz, Germany, and
Vernadsky Institute of Geochemistry, Moscow 117975, Russia

Dmitry V. Kuzmin

Max-Planck-Institut für Chemie, Postfach 3060, 55020 Mainz, Germany, and
Institute of Geology and Mineralogy SB RAS, Novosibirsk 630090, Russia

Rainer Thomas

GeoForschungsZentrum Potsdam, Telegrafenberg B 120, D-14473 Potsdam, Germany

ABSTRACT

The volcanic hazard potential of Mount Etna volcano is currently nourished by long-lasting, powerful eruptions of basaltic magmas coupled with increased seismicity and ground deformation, and the world's largest discharge of volcanic gases. The current evolutionary cycle of Mount Etna activity is consistent with subduction-related chemical modifications of the mantle source. Arrival of a new mantle-derived magma batch beneath the volcano has been hypothesized, but is still elusive among the erupted products. Here we demonstrate petrological and geochemical affinities between the magmas supplying modern eruptions and high-Mg, fall-stratified (FS) basalts ejected violently ~4 k.y. ago. The FS primitive magmas (~13 wt% MgO) are characteristically volatile enriched (at least 3.8 wt% H₂O and 3300 ppm CO₂), and bear a trace element signature of a garnet-bearing, metasomatized source (high Gd/Yb, K/La, U/Nb, Pb/Ce, Ca/Al). They started crystallizing olivine (Fo₉₁), clinopyroxene (Mg# 92.5), and Cr spinel deep in the plumbing system (>5 kbar), contributing to the cumulate piles at depth and to differentiated alkaline basalt and trachybasalt magmas in the shallow conduit. Continuous influx of mantle-derived, volatile-rich magmas, such as those that supplied the FS fallout, provides a good explanation for major compositional and eruptive features of Mount Etna.

Keywords: Mount Etna, volcano, picrite, melt inclusions, volatiles, degassing, mantle, metasomatism.

INTRODUCTION

Mount Etna, positioned at the destructive margin marking the complex collision zone between the African and European plates (Gueguen et al., 1998), is the largest volcano in Europe and one of the most active in the world. The present edifice of Mount Etna was constructed with fractionation products of mantle-derived magmas, starting ~0.5 m.y. ago with the eruption of tholeiitic lavas, followed by transitional lavas, eventually trending to SiO₂-undersaturated, alkaline basalts with distinct trace element and isotope signatures (Armienti et al., 2004; Corsaro and Pompilio, 2004; Tanguy et al., 1997). The compositional changes of the Mount Etna magmatic system were temporally and genetically linked

to continuous modifications in the mantle-lithospheric source(s), caused by metasomatic enrichment in volatiles and incompatible trace elements, and melting-related depletion in basaltic components (Kamenetsky and Clocchiatti, 1996). Dramatic changes in both the source composition and the volcano's eruptive dynamics have been tentatively ascribed to modification of the adjacent subduction system (Schiano et al., 2001; Tonarini et al., 2001), driven by vertical asthenospheric flow in response to the rollback of the subducting Ionian slab (Dogliani et al., 2001; Gvirtzman and Nur, 1999).

From a historical perspective, Mount Etna shows increasing activity in recent time; with the frequency of explosive (doubled since the fall-stratified [FS] eruption of high-magnesian magmas ~4 k.y. ago) and effusive eruptions has

increased significantly (Coltelli et al., 2000). Since the early 1970s, the magmatic output of Mount Etna has increased by a factor of 4, and more than 120 eruptions have been recorded in the past 20 yr (Branca and Del Carlo, 2005). Notably, the erupted magmas have become more potassic and richer in ⁸⁷Sr and ¹¹B (Armienti et al., 2004; Clocchiatti et al., 1988; Tanguy et al., 1997; Tonarini et al., 2001), and have been accompanied by enormous output of volcanic gases CO₂, SO₂, and halogens (Allard, 1997; Allard et al., 1991; D'Alessandro et al., 1997). Although the present-day magmas are increasingly magnesian and volatile rich (Clocchiatti et al., 2004; Corsaro and Pompilio, 2004; Métrich et al., 2004; Spilliaert et al., 2006), and their radiogenic and stable isotope compositions require a mantle origin (Allard et al., 1991;

*E-mail: Dima.Kamenetsky@utas.edu.au.

Armienti et al., 2004; D'Alessandro et al., 1997; Marty et al., 1994; Tonarini et al., 2001), their MgO (<8.2 wt%) and Ni (<50 ppm) contents, and forsterite contents of olivine phenocrysts (<84 mol% Fo), are too low to have been in equilibrium with a typical peridotite mantle.

However, the indications of a primitive melt entering the plumbing system of Mount Etna from a mantle reservoir are explicit. In addition to the isotope systematics, these include records of large piles of mafic cumulates extending to depths of 18 km beneath the volcano (De Gori et al., 2005) and the excessive (compared to amounts degassed from the erupted magmas) supply of volatiles to the volcanic clouds (Allard et al., 1991; D'Alessandro et al., 1997). As primitive magmas have rarely been ejected in the eruptive history of Mount Etna (Corsaro and Pompilio, 2004) and have not reached the surface since the FS eruption, we used the latter to elucidate controls on recent and current volcanic activity.

FS ERUPTION: ROCKS, MINERALS, AND MELT INCLUSIONS

The 3930 ± 60 B.P. (^{14}C age; Coltelli et al., 2000) FS eruption produced a pyroclastic fall-out deposit of 0.183 km^3 that covered the eastern flanks of Mount Etna (Coltelli et al., 2005). The deposit reaches 110 cm in thickness at ~ 7 km from the summit craters, and its thickness and dispersal indicate that the eruption column reached significant heights (18–20 km; Coltelli et al., 2005), and thus had high explosive (sub-Plinian) energy. The erupted products are a highly vesiculated (30–60 vol%) scoria lapilli containing euhedral crystals of olivine (<8%), clinopyroxene (<3%), and Cr spinel (<0.1%). Despite relatively low amounts of ferromagnesian phenocrysts, the rocks are high-Mg basalt to picrite in composition (12–17 wt% MgO; GSA Data Repository Table DR1¹) with high Ni and Cr abundances (135–285 and 920–1330 ppm, respectively), characteristic of primitive unfractionated magmas. The rocks are extremely enriched in CaO relative to Al_2O_3 ($\text{CaO}/\text{Al}_2\text{O}_3 = 1.1\text{--}1.2$), despite the low modal abundances of clinopyroxene phenocrysts. The FS rocks are depleted in heavy rare earth elements (HREE) and high-field strength elements (HFSE) relative to REEs of higher and similar incompatibility, respectively (Fig. 1). They are also relatively enriched in large-ion lithophile

¹GSA Data Repository item 2007056, methods and calculations, and Tables DR1 (major and trace element compositions of FS rocks and olivine-hosted melt inclusions), Table DR2 (representative compositions of clinopyroxene inclusions and host olivine from FS rocks), and Table DR3 (representative compositions of Cr-spinel inclusions and host olivine from FS rocks), is available online at www.geosociety.org/pubs/ft2007.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

elements (LILE—K and Sr) and Pb (Fig. 1). Depletions and enrichments demonstrated by the ratios of similarly incompatible elements (e.g., K/La, Nb/U, Ce/Pb, Ti/Gd) are typical of subduction-related magmas and commonly

attributed to selective addition of LILE > light REE > HFSE to a mantle peridotite by melts and/or fluids derived from a subducting crust. High Gd/Yb in the magmas can be taken as evidence of residual garnet in the source.




Figure 1. Primitive mantle normalized (after Sun and McDonough, 1989) trace element compositions of fall-stratified (FS) rocks (open circles, average of 6 samples), melt inclusions hosted in FS olivine phenocrysts (closed circles, average of 23 melt inclusions; bars show one standard deviation), and Mount Etna lavas erupted in 2001–2002 (diamonds, average of 61 samples; after Andronico et al., 2005; Corsaro et al., 2006).




Figure 2. Comparison between compositions of fall-stratified (FS) parental magmas (exemplified by recalculated melt inclusions), FS whole rocks, and modern lavas of Mount Etna (Andronico et al., 2005; Corsaro et al., 2006). Note significant variations in trace elements among individual melt inclusions. Symbols as in Figure 1.

High-Mg compositions of the FS olivine phenocrysts (Fo 90–91 mol% in the majority of crystals) and clinopyroxene (Mg# 96–92.5) also point to the primitive nature of their parental magmas (see Tables DR2 and DR3). Early crystallization of FS clinopyroxene, confirmed by the occurrence of clinopyroxene inclusions in most forsteritic olivine, may reflect high Ca abundance in the parental magma and high-pressure crystallization conditions (Grove et al., 1992). Spinel microphenocrysts and inclusions in olivine have extremely high Cr (Cr# 70–80 mol%) and moderately low Ti (0.6–1.0 wt% TiO₂) compositions that are also typical of many primitive alkaline island arc magmas, including those from nearby Vulcano, Aeolian arc (Kamenetsky and Clocchiatti, 1996; Kamenetsky et al., 2001).

Olivine-hosted melt inclusions, analyzed for major, trace, and volatile elements, using different microbeam techniques, and subsequently corrected for postentrapment modifications (see Data Repository [see footnote 1] for the details of analytical methods and calculations) provide more insight into the nature of the FS parental magmas (Figs. 2 and 3). Melt inclusions (as large as 250 μm) have negative-crystal shape with scalloped edges and appear as brownish glass with a spherical vapor bubble (Fig. 3), which is sometimes larger than typical shrinkage bubbles (i.e., >5 vol%). The bubble walls are invariably lined with tiny Ca-Mg carbonate crystals (Fig. 3).

Melt inclusion compositions show a limited range in MgO (12.6 ± 0.3 wt%) and match the bulk rock major and trace element compositions (Figs. 1 and 2). Thus, the melt inclusions support our conclusions about high-Mg and high-Ca compositions of the FS parental melt with the trace element systematics reminiscent of subduction-related magmatism. Although the average composition of melt inclusions is indistinguishable from the average bulk rock composition, the observed chemical variability among individual inclusions, especially in trace elements (Figs. 1 and 2), is very large and cannot have been induced by crystallization because the compositions of host olivines are similar.

Melt inclusions have moderately high Cl and S (average 0.15 and 0.17 wt%, respectively), but are exceptionally enriched in H₂O and CO₂ for basaltic melts, in general, and primitive high-Mg basalt, in particular (Fig. 3). The original CO₂ content of trapped melts was undoubtedly higher than measured in glasses, because carbon present in the bubbles (gaseous phase and carbonates) is not quantified. Because H₂O does not correlate with elements of similar incompatibility (see large variability in trace elements in Fig. 2), the effects of degassing cannot be excluded. Nevertheless, the highest measured abundances of dissolved volatiles indicate very




Figure 3. Relationships between H₂O and CO₂ contents in melt inclusions from fall-stratified (FS) olivine (circles), phenocrysts in recent Mount Etna lavas (diamonds; after Métrich et al., 2004; Spilliaert et al., 2006), and arc basalts (dotted outline; after Wallace, 2005). Note that adequate comparison of data from different sources requires normalization to same degree of fractionation. Isobars are calculated at 1220 °C and 47.5 wt% SiO₂ in melt using Volatilecalc (Newman and Lowenstern, 2002). Inset shows typical H₂O- and CO₂-rich silicate melt inclusion with CO₂-vapor bubble in FS olivine. Bubble walls are coated by tiny Ca-Mg-bearing carbonates (determined by laser Raman spectroscopy, GeoForschungsZentrum Potsdam, Germany), thus initial CO₂ content was even higher than measured.

deep crystallization (at least 400–500 MPa, Fig. 3) of clinopyroxene and olivine from the FS parental melt.

IMPLICATIONS FOR MODERN ERUPTIONS

The evolution of the FS parental melt upon decompression, cooling, degassing, and extensive (50%–60%) crystallization of olivine and clinopyroxene (with the plagioclase liquidus depressed by high H₂O contents) can hypothetically account for the compositions of modern lavas at Mount Etna. The latter show the compositional trends in major and trace elements that can be tracked back to the FS compositions (Fig. 2), the geochemical signature consistent with their derivation from the high-magnesian, FS-type magma (Fig. 1). Water and CO₂ contents in the melt inclusions of modern eruptions (Del Carlo and Pompilio, 2004; Métrich et al., 2004; Métrich and Rutherford, 1998; Spilliaert et al., 2006) also point to the volatile-enriched character of parental magmas (Fig. 3); however, linking the FS magmas and the present-day magmas requires a more comprehensive interpretation of the behavior of volatile elements. Quantitative relationships between the FS and modern magmas are hampered by current lack of relevant experimental data to model the olivine-clinopyroxene cotectic, and undefined extents of primitive magma replenishment, interaction

with earlier cumulates, mixing with evolved magmas in the conduit, and degassing.

In historic time, and most evident recently, Mount Etna showed prominent geochemical changes in the erupted products (Armienti et al., 2004; Clocchiatti et al., 1988; Tanguy et al., 1997; Tonarini et al., 2001) and dramatically increased volcanic activity, seismicity, and ground deformation (Branca and Del Carlo, 2005; Patane et al., 2003). Uncommon elsewhere in the world, frequent explosive eruptions of basaltic magmas in the latest volcanic cycles of Mount Etna are coupled with the largest (for a single volcano) emissions of volcanic gases CO₂ and SO₂ (Allard, 1997; Allard et al., 1991; D'Alessandro et al., 1997). Another unusual aspect of the Mt. Etna magmatic system is the vertical extent (up to 18 km) of the magma storage region and related mafic cumulates indicated by geophysical and geochemical data (Allard, 1997; Chiarabba et al., 2000; De Gori et al., 2005; Laigle et al., 2000; Tanguy et al., 1997). A prerequisite to all features that make present-day Mount Etna an extraordinary magmatic system is arrival of new primitive magma into the feeding system. Such magma, hypothesized in several works (Clocchiatti et al., 2004; Métrich et al., 2004; Patane et al., 2003; Spilliaert et al., 2006) but not sampled directly among recent volcanics, can be best approximated by an unevolved, mantle-derived,

FS-type magma, established in our study. The FS-type melt demonstrates properties critical for positive magma buoyancy, i.e., originally high abundances of dissolved H₂O and CO₂ that were further increased by crystallization of olivine and clinopyroxene at depth. Subsequent decompression promoted early saturation of the melt in volatiles, vesiculation, and rapid ascent of the low-density, highly buoyant melt-bubble mixture. Thus, the explosive dynamics of modern Mount Etna can be related to a deep supply of volatile-rich, high-Mg magmas. The recognition of such magmas in the plumbing system of Mount Etna is also important in linking shallow-crust and surface processes to the evolution of the mantle source region. The FS-type melt provides compelling compositional evidence that Mount Etna is currently supplied from a source, increasingly fluxed by melts and fluids, in a continuously changing tectonic framework.

ACKNOWLEDGMENTS

This paper has greatly benefited from comments by Malcolm Rutherford, Paul Wallace, and Dawnika L. Blatter. This research was supported by a Research Fellowship and Discovery grant (Australian Research Council, Australia) and F.W. Bessel Award (A. von Humboldt Foundation, Germany) to V. Kamenetsky, and an Istituto Nazionale di Geofisica e Vulcanologia–Dipartimento della Protezione Civile grant to M. Pompilio.

REFERENCES CITED

- Allard, P., 1997, Endogenous magma degassing and storage at Mount Etna: *Geophysical Research Letters*, v. 24, p. 2219–2222, doi: 10.1029/97GL021011.
- Allard, P., Carbonnelle, J., Dajlevic, D., Lebronec, J., Morel, P., Robe, M.C., Maurenas, J.M., Faivrepierrre, R., Martin, D., Sabroux, J.C., and Zettwoog, P., 1991, Eruptive and diffuse emissions of CO₂ from Mount Etna: *Nature*, v. 351, p. 387–391, doi: 10.1038/351387a0.
- Andronico, D., Branca, S., Calvari, S., Burton, M., Caltabiano, T., Corsaro, R.A., Del Carlo, P., Garfi, G., Lodato, L., Miraglia, L., Mure, F., Neri, M., Pecora, E., Pompilio, M., Salerno, G., and Spampinato, L., 2005, A multi-disciplinary study of the 2002–03 Etna eruption: Insights into a complex plumbing system: *Bulletin of Volcanology*, v. 67, p. 314–330, doi: 10.1007/s00445-004-0372-8.
- Armienti, P., Tonarini, S., D’Orazio, M., and Innocenti, F., 2004, Genesis and evolution of Mt. Etna alkaline lavas: Petrological and Sr-Nd-B isotope constraints: *Periodico di Mineralogia*, v. 73, p. 29–52.
- Branca, S., and Del Carlo, P., 2005, Types of eruptions of Etna volcano AD 1670–2003: Implications for short-term eruptive behaviour: *Bulletin of Volcanology*, v. 67, p. 732–742, doi: 10.1007/s00445-005-0412-z.
- Chiarabba, C., Amato, A., Boschi, E., and Barberi, F., 2000, Recent seismicity and tomographic modeling of the Mount Etna plumbing system: *Journal of Geophysical Research*, v. 105, p. 10,923–10,938, doi: 10.1029/1999JB900427.
- Clocchiatti, R., Joron, J.L., and Treuil, M., 1988, The role of selective alkali contamination in the evolution of recent historic lavas of Mt. Etna: *Journal of Volcanology and Geothermal Research*, v. 34, p. 241–249, doi: 10.1016/0377-0273(88)90036-4.
- Clocchiatti, R., Condomines, M., Guenet, N., and Tanguy, J.C., 2004, Magma changes at Mount Etna: The 2001 and 2002–2003 eruptions: *Earth and Planetary Science Letters*, v. 226, p. 397–414, doi: 10.1016/j.epsl.2004.07.039.
- Coltelli, M., Del Carlo, P., and Vezzoli, L., 2000, Stratigraphic constraints for explosive activity in the past 100 ka at Etna Volcano, Italy: *International Journal of Earth Sciences*, v. 89, p. 665–677, doi: 10.1007/s005310000117.
- Coltelli, M., Del Carlo, P., Pompilio, M., and Vezzoli, L., 2005, Explosive eruption of a picrite: The 3930 BP subplinian eruption of Etna volcano (Italy): *Geophysical Research Letters*, v. 32, L23307, doi: 10.1029/2005GL024271.
- Corsaro, R.A., and Pompilio, M., 2004, Dynamics of magma at Mount Etna, in Bonaccorso, A., et al., eds., *Mt Etna: Volcano Laboratory*, Volume 143: Washington, D.C., American Geophysical Union, p. 91–110.
- Corsaro, R.A., Miraglia, L., and Pompilio, M., 2006, Petrologic evidence of a complex plumbing system feeding the July–August 2001 eruption of Mt. Etna, Sicily, Italy: *Bulletin of Volcanology*, doi: 10.1007/s00445-006-0083-4.
- D’Alessandro, W., Giammanco, S., Parello, F., and Valenza, M., 1997, CO₂ output and δ¹³C(CO₂) from Mount Etna as indicators of degassing of shallow asthenosphere: *Bulletin of Volcanology*, v. 58, p. 455–458, doi: 10.1007/s004450050154.
- De Gori, P., Chiarabba, C., and Patane, D., 2005, *Qp* structure of Mount Etna: Constraints for the physics of the plumbing system: *Journal of Geophysical Research*, v. 110, p. B05303, doi: 10.1029/2003JB002875.
- Del Carlo, P., and Pompilio, M., 2004, The relationship between volatile content and the eruptive style of basaltic magma: The Etna case: *Annals of Geophysics*, v. 47, p. 1423–1432.
- Dogliani, C., Innocenti, F., and Mariotti, G., 2001, Why Mt Etna?: *Terra Nova*, v. 13, p. 25–31, doi: 10.1046/j.1365-3121.2001.00301.x.
- Grove, T.L., Kinzler, R.J., and Bryan, W.B., 1992, Fractionation of mid-ocean ridge basalt (MORB), in Phipps Morgan, J., et al., eds., *Mantle flow and melt generation at mid-ocean ridges*: American Geophysical Union Geophysical Monograph 71, p. 281–310.
- Gueguen, E., Dogliani, C., and Fernandez, M., 1998, On the post-25 Ma geodynamic evolution of the western Mediterranean: *Tectonophysics*, v. 298, p. 259–269, doi: 10.1016/S0040-1951(98)00189-9.
- Gvirtzman, Z., and Nur, A., 1999, The formation of Mount Etna as the consequence of slab rollback: *Nature*, v. 401, p. 782–785, doi: 10.1038/44555.
- Kamenetsky, V., and Clocchiatti, R., 1996, Primitive magmatism of Mt. Etna: Insights from mineralogy and melt inclusions: *Earth and Planetary Science Letters*, v. 142, p. 553–572, doi: 10.1016/0012-821X(96)00115-X.
- Kamenetsky, V.S., Crawford, A.J., and Meffre, S., 2001, Factors controlling chemistry of magmatic spinel: An empirical study of associated olivine, Cr-spinel and melt inclusions from primitive rocks: *Journal of Petrology*, v. 42, p. 655–671, doi: 10.1093/petrology/42.4.655.
- Laigle, M., Hirn, A., Sapin, M., Lepine, J.C., Diaz, J., Gallart, J., and Nicolich, R., 2000, Mount Etna dense array local earthquake P and S tomography and implications for volcanic plumbing: *Journal of Geophysical Research*, v. 105, p. 21,633–21,646, doi: 10.1029/2000JB900190.
- Marty, B., Trull, T., Lussiez, P., Basile, I., and Tanguy, J.C., 1994, He, Ar, O, Sr and Nd isotope constraints on the origin and evolution of Mount Etna magmatism: *Earth and Planetary Science Letters*, v. 126, p. 23–39, doi: 10.1016/0012-821X(94)90240-2.
- Métrich, N., and Rutherford, M.J., 1998, Low pressure crystallization paths of H₂O-saturated basaltic-hawaiitic melts from Mt Etna: Implications for open-system degassing of basaltic volcanoes: *Geochimica et Cosmochimica Acta*, v. 62, p. 1195–1205, doi: 10.1016/S0016-7037(98)00048-9.
- Métrich, N., Allard, P., Spilliaert, N., Andronico, D., and Burton, M., 2004, 2001 flank eruption of the alkali- and volatile-rich primitive basalt responsible for Mount Etna’s evolution in the last three decades: *Earth and Planetary Science Letters*, v. 228, p. 1–17, doi: 10.1016/j.epsl.2004.09.036.
- Newman, S., and Lowenstern, J.B., 2002, Volatile-Calc: A silicate melt-H₂O-CO₂ solution model written in Visual Basic for excel: *Computers & Geosciences*, v. 28, p. 597–604, doi: 10.1016/S0098-3004(01)00081-4.
- Patane, D., De Gori, P., Chiarabba, C., and Bonaccorso, A., 2003, Magma ascent and the pressurization of Mount Etna’s volcanic system: *Science*, v. 299, p. 2061–2063, doi: 10.1126/science.1080653.
- Schiano, P., Clocchiatti, R., Ottolini, L., and Busa, T., 2001, Transition of Mount Etna lavas from a mantle-plume to an island-arc magmatic source: *Nature*, v. 412, p. 900–904, doi: 10.1038/35091056.
- Spilliaert, N., Allard, P., Métrich, N., and Sobolev, A.V., 2006, Melt inclusion record of the conditions of ascent, degassing and extrusion of volatile-rich alkali basalt during the powerful 2002 flank eruption of Mount Etna (Italy): *Journal of Geophysical Research*, v. 111, p. B04203, doi: 10.1029/2005JB003934.
- Sun, S.-S., and McDonough, W.F., 1989, Chemical and isotopic systematics of oceanic basalts: Implications for mantle composition and processes, in Saunders, A.D., and Norry, M.J., eds., *Magmatism in the ocean basins*: Geological Society [London] Special Publication 42, p. 313–345.
- Tanguy, J.C., Condomines, M., and Kieffer, G., 1997, Evolution of the Mount Etna magma: Constraints on the present feeding system and eruptive mechanism: *Journal of Volcanology and Geothermal Research*, v. 75, p. 221–250, doi: 10.1016/S0377-0273(96)00065-0.
- Tonarini, S., Armienti, P., D’Orazio, M., and Innocenti, F., 2001, Subduction-like fluids in the genesis of Mt. Etna magmas: Evidence from boron isotopes and fluid mobile elements: *Earth and Planetary Science Letters*, v. 192, p. 471–483, doi: 10.1016/S0012-821X(01)00487-3.
- Wallace, P.J., 2005, Volatiles in subduction zone magmas: Concentrations and fluxes based on melt inclusion and volcanic gas data: *Journal of Volcanology and Geothermal Research*, v. 140, p. 217–240, doi: 10.1016/j.jvolgeores.2004.07.023.

Manuscript received 4 July 2006
 Revised manuscript received 30 October
 Manuscript accepted 31 October 2006

Printed in USA